



Acer rubrum Red Maple¹

Edward F. Gilman and Dennis G. Watson²

INTRODUCTION

Red Maple has an oval shape and is a fast grower with strong wood, reaching a height of 75 feet (Fig. 1). Unless irrigated or on a wet site, Red Maple is best used north of USDA hardiness zone 9. Trees are often much shorter in the southern part of its range unless growing next to a stream or on a wet site. This tree is preferred over Silver Maple or Boxelder when a fast growing Maple is needed. When planting the species *Acer rubrum*, select only those which have been grown from seed sources in your area. The newly emerging leaves and red flowers and fruits signal that spring has come. They appear in December and January in Florida, later in the northern part of its range. The seeds of red maple are quite popular with squirrels and birds. This tree is sometimes confused with red-leaved cultivars of Norway Maple.

GENERAL INFORMATION

Scientific name: *Acer rubrum*

Pronunciation: AY-ser ROO-brum

Common name(s): Red Maple, Swamp Maple

Family: *Aceraceae*

USDA hardiness zones: 4 through 9 (Fig. 2)

Origin: native to North America

Uses: Bonsai; wide tree lawns (>6 feet wide); medium-sized tree lawns (4-6 feet wide); recommended for buffer strips around parking lots or for median strip plantings in the highway; near a deck or patio; reclamation plant; screen; shade tree; residential street tree

Availability: generally available in many areas within its hardiness range




Figure 1. Mature Red Maple.

DESCRIPTION

Height: 60 to 75 feet

Spread: 25 to 35 feet

Crown uniformity: irregular outline or silhouette

Crown shape: oval; round; upright

Crown density: moderate

Growth rate: fast

Texture: medium

1. This document is adapted from Fact Sheet ST-41, a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Publication date: November 1993.
2. Edward F. Gilman, associate professor, Environmental Horticulture Department; Dennis G. Watson, associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.



Figure 2. Shaded area represents potential planting range.

Foliage

Leaf arrangement: opposite/subopposite (Fig. 3)

Leaf type: simple

Leaf margin: lobed; incised; serrate

Leaf shape: ovate

Leaf venation: palmate

Leaf type and persistence: deciduous

Leaf blade length: 2 to 4 inches

Leaf color: green

Fall color: orange; red; yellow

Fall characteristic: showy

Flower

Flower color: red

Flower characteristics: showy; spring flowering;
winter flowering

Fruit

Fruit shape: elongated

Fruit length: 1 to 3 inches

Fruit covering: dry or hard

Fruit color: red

Fruit characteristics: attracts birds; attracts squirrels and other mammals; no significant litter problem; showy

Trunk and Branches

Trunk/bark/branches: bark is thin and easily damaged from mechanical impact; droop as the tree grows, and will require pruning for vehicular or pedestrian clearance beneath the canopy; not particularly showy; should be grown with a single leader; no thorns

Pruning requirement: requires pruning to develop strong structure

Breakage: susceptible to breakage either at the crotch due to poor collar formation, or the wood itself is weak and tends to break

Current year twig color: gray; reddish

Current year twig thickness: medium

Wood specific gravity: 0.54




Figure 3. Foliage of Red Maple.

Culture

Light requirement: tree grows in part shade/part sun;
tree grows in full sun

Soil tolerances: clay; loam; sand; acidic; extended
flooding; well-drained

Drought tolerance: moderate

Aerosol salt tolerance: low

Soil salt tolerance: poor

Other

Roots: surface roots can lift sidewalks or interfere
with mowing

Winter interest: tree has winter interest due to
unusual form, nice persistent fruits, showy winter
trunk, or winter flowers

Outstanding tree: not particularly outstanding

Invasive potential: little, if any, potential at this time

Ozone sensitivity: tolerant

Verticillium wilt susceptibility: susceptible

Pest resistance: long-term health usually not
affected by pests

USE AND MANAGEMENT

The outstanding ornamental characteristic of Red Maple is red, orange or yellow fall color (sometimes on the same tree) lasting several weeks. Red maple is often one of the first trees to color up in autumn, and it puts on one of the most brilliant displays of any tree but trees vary greatly in fall color and intensity. Cultivars are more consistently colored.

The tree makes the best growth in wet places and has no other particular soil preference except chlorosis may develop on alkaline soil where it also grows poorly. It is well-suited as a street tree in northern and mid-south climates in residential and other suburban areas but the bark is thin and easily damaged by mowers. Irrigation is often needed to support street tree plantings in well-drained soil in the south. Roots can raise sidewalks as silver maples can but they have a less aggressive root system and so they make a good street tree. Surface roots beneath the canopy can make mowing difficult.

Red Maple is easily transplanted and usually develops surface roots in soil ranging from well-drained sand to clay. It is not especially drought-tolerant, particularly in the southern part of the range, but selected individual trees can be found growing on dry sites. This trait shows the wide range of genetic diversity in the species. Branches often grow upright through the crown forming poor attachments to the trunk. These should be removed in the nursery or after planting in the landscape to help prevent branch failure in older trees during storms. Select branches with a wide angle from the trunk and prevent branches from growing larger than half the diameter of the trunk.

A number of cultivars are listed. Due to graft-incompatibility problems which cause the tree to break apart, preference should be given to cultivars produced on their own roots. In the northern and southern end of the range, choose cultivars with regional adaptation. The cultivars are: 'Armstrong' - upright growth habit, almost columnar, somewhat prone to splitting branches due to tight crotches, 50 feet tall; 'Autumn Flame' - 45 feet tall, round, above average fall color; 'Bowhall' - upright growth habit, branches form embedded bark, graft incompatibility on grafted trees; 'Gerling' - densely branched, broadly pyramidal, about 35 feet tall when mature; 'October Glory' - above average fall color, excellent tree, retains leaves late, 60 feet tall; 'Red Sunset' - above average orange to red fall color, does well in the south in USDA hardiness zone 8,

probably the best cultivar for the deep south, oval, 50 feet tall; 'Scanlon' - upright growth habit; 'Schlesinger' - good fall color, rapid growth rate; 'Tilford' - globe-shaped crown. Variety *drummondii* suitable in USDA hardiness zone 8.

There is a recently introduced hybrid cross between Red and Silver Maple called Hybrid Maple (*Acer x fremanii*). Cultivars of this hybrid include 'Armstrong' with a narrow columnar crown to 35 feet tall, 'Autumn Blaze' with an oval crown to 50 feet tall, 'Celebration' with a narrow upright crown and a strong central leader to 50 or 60 feet tall, 'Celzam' with a narrow oval crown to 50 feet tall, and 'Scarlet Sentenial' with great fall color, and oval crown to 40 feet tall. The culture of these trees is probably similar to Red Maple.

Pests

Leaf stalk borer and petiole-borer cause the same type of injury. Both insects bore into the leaf stalk just below the leaf blade. The leaf stalk shrivels, turns black, and the leaf blade falls off. The leaf drop may appear heavy but serious injury to a healthy tree is rare.

Gall mites stimulate the formation of growths or galls on the leaves. The galls are small but can be so numerous that individual leaves curl up. The most common gall is bladder gall mite found on silver maple. The galls are round and at first green but later turn red, then black, then dry up. Galls of other shapes are seen less frequently on other types of maples. Galls are not serious, so chemical controls are not needed.

Crimson erineum mite is usually found on silver maple and causes the formation of red fuzzy patches on the lower leaf surfaces. The problem is not serious so control measures are not suggested.

Aphids infest maples, usually Norway Maple, and may be numerous at times. High populations can cause leaf drop. Another sign of heavy aphid infestation is honey dew on lower leaves and objects beneath the tree. Aphids are controlled by spraying or they may be left alone. If not sprayed, predatory insects will bring the aphid population under control.

Scales are an occasional problem on maples. Perhaps the most common is cottony maple scale. The insect forms a cottony mass on the lower sides of branches. Scales are usually controlled with

horticultural oil sprays. Scales may also be controlled with well-timed sprays to kill the crawlers.

If borers become a problem it is an indication the tree is not growing well. Controlling borers involves keeping trees healthy. Chemical controls of existing infestations are more difficult. Proper control involves identification of the borer infesting the tree then applying insecticides at the proper time.

Twig borers can cause die-back of the terminal 8 to 12 inches of small-diameter branches. This is usually not serious and does not require control measures, but it can be a problem on young trees in the nursery.

Diseases

Anthracnose is more of a problem in rainy seasons. The disease resembles, and may be confused with, a physiological problem called scorch. The disease causes light brown or tan areas on the leaves. Anthracnose may be controlled by fungicides sprayed on as leaves open in the spring. Two additional sprays at two-week intervals will be needed. The disease is most common on sugar and silver maples and boxelder. Other maples may not be affected as severely. Sprays may need to be applied by a commercial applicator having proper spray equipment.

Girdling roots grow around the base of the trunk rather than growing away from it. As both root and trunk increase in size, the root chokes the trunk. Girdling roots are detected by examining the base of the trunk. The lack of trunk flare at ground level is a symptom. The portion of the trunk above a girdling root does not grow as rapidly as the rest so may be slightly depressed. The offending root may be on the surface or may be just below the sod. The tree crown shows premature fall coloration and death of parts of the tree in more serious cases. If large portions of the tree have died it may not be worth saving. Girdling roots are functional roots so when removed a portion of the tree may die. When the girdling root is large the treatment is as harmful as the problem. After root removal, follow-up treatment includes watering during dry weather. The best treatment for girdling roots is prevention by removing or cutting circling roots at planting or as soon as they are detected on young trees.

Scorch may occur during periods of high temperatures accompanied by wind. Trees with diseased or inadequate root systems will also show

scorching. When trees do not get enough water they scorch. Scorch symptoms are light brown or tan dead areas between leaf veins. The symptoms are on all parts of the tree or only on the side exposed to sun and wind. Scorching due to dry soil may be overcome by watering. If scorching is due to an inadequate or diseased root system, watering may have no effect.

Nutrient deficiency symptoms are yellow or yellowish-green leaves with darker green veins. The most commonly deficient nutrient on maple is manganese. Implanting capsules containing a manganese source in the trunk will alleviate the symptoms. Test soil samples to determine if the soil pH is too high for best manganese availability. Plants exposed to weed killers may also show similar symptoms.

Tar spot and a variety of leaf spots cause some concern among homeowners but are rarely serious enough for control.