



***Acer campestre* 'Evelyn' 'Evelyn' Hedge Maple¹**

Edward F. Gilman and Dennis G. Watson²

INTRODUCTION

'Evelyn' Hedge Maple is usually low-branched with an upright-rounded form (Fig. 1). It may not be as cold hardy as the species. This cultivar of hedge maple has a more consistent upright branching habit which probably makes it more suited for street and parking lot planting than the species. The branches are slender and branch profusely, lending a fine texture to the landscape particularly during winter. Little pruning would be needed to keep the lower crown clear from vehicle and pedestrians since branches grow upright. The tree eventually reaches a height and spread of 30 to 35 feet but it grows slowly. The small stature and vigorous growth make this an excellent street tree for residential areas, or perhaps in downtown urban sites. However, it grows a little too tall for planting beneath some power lines. It is also suitable as a patio or yard shade tree because it stays small and creates dense shade.

GENERAL INFORMATION

Scientific name: *Acer campestre* 'Evelyn'

Pronunciation: AY-ser kam-PESS-tree

Common name(s): 'Evelyn' Hedge Maple

Family: *Aceraceae*

USDA hardiness zones: 6 through 8A (Fig. 2)

Origin: not native to North America

Uses: Bonsai; hedge; large parking lot islands (> 200 square feet in size); wide tree lawns (>6 feet wide); medium-sized parking lot islands (100-200 square feet in size); medium-sized tree lawns (4-6 feet wide); recommended for buffer strips around parking lots or for median strip plantings in the highway; near a deck




Figure 1. Middle-aged 'Evelyn' Hedge Maple.

or patio; screen; trainable as a standard; shade tree; small parking lot islands (< 100 square feet in size); narrow tree lawns (3-4 feet wide); specimen; sidewalk cutout (tree pit); residential street tree; tree has been successfully grown in urban areas where air pollution,

1. This document is adapted from Fact Sheet ST-12, a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Publication date: November 1993.
2. Edward F. Gilman, associate professor, Environmental Horticulture Department; Dennis G. Watson, associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.



Figure 2. Shaded area represents potential planting range.

poor drainage, compacted soil, and/or drought are common

Availability: grown in small quantities by a small number of nurseries

DESCRIPTION

Height: 30 to 35 feet

Spread: 30 to 35 feet

Crown uniformity: symmetrical canopy with a regular (or smooth) outline, and individuals have more or less identical crown forms

Crown shape: upright

Crown density: dense

Growth rate: medium

Texture: fine

Foliage

Leaf arrangement: opposite/subopposite (Fig. 3)

Leaf type: simple

Leaf margin: lobed; entire

Leaf shape: star-shaped

Leaf venation: palmate

Leaf type and persistence: deciduous

Leaf blade length: 2 to 4 inches

Leaf color: green

Fall color: yellow

Fall characteristic: showy

Flower

Flower color: green

Flower characteristics: inconspicuous and not showy; spring flowering

Fruit

Fruit shape: oval

Fruit length: 1 to 3 inches

Fruit covering: dry or hard

Fruit color: brown; green

Fruit characteristics: does not attract wildlife; inconspicuous and not showy; no significant litter problem

Trunk and Branches

Trunk/bark/branches: bark is thin and easily damaged from mechanical impact; grow mostly upright and will not droop; not particularly showy; should be grown with a single leader; no thorns




Figure 3. Foliage of 'Evelyn' Hedge Maple.

Pruning requirement: requires pruning to develop strong structure

Breakage: resistant

Current year twig color: brown

Current year twig thickness: medium; thin

Culture

Light requirement: tree grows in part shade/part sun; tree grows in full sun

Soil tolerances: clay; loam; sand; acidic; alkaline; well-drained

Drought tolerance: high

Aerosol salt tolerance: moderate

Other

Roots: surface roots are usually not a problem

Winter interest: tree has winter interest due to unusual form, nice persistent fruits, showy winter trunk, or winter flowers

Outstanding tree: tree has outstanding ornamental features and could be planted more

Invasive potential: little, if any, potential at this time

Verticillium wilt susceptibility: susceptible

Pest resistance: no pests are normally seen on the tree

USE AND MANAGEMENT

The tree excels in its ability to tolerate dry, alkaline soil but some protection from open winds is helpful. Not for highly compacted soil. It is well-suited for and looks great during drought in a partially shaded location or on the north side of a building. The common name alludes to the plants tolerance of severe pruning, and it will make a dense, tall screen, whether pruned or not. Branches are arranged closely on the trunk and some pruning is usually desirable to create a well-formed tree. The main ornamental feature is the consistently bright yellow fall color. Prune early in the life of the tree to develop several major branches well-spaced along a central trunk. This will improve the durability of the tree compared to trees with many upright and spreading branches originating from one point on the trunk.

The cultivar 'Compactum' is dwarf, 'Postelense' has golden leaves.

Pests

Pests are usually not a problem but there are some potential problems.

Leaf stalk borer and petiole-borer cause the same type of injury. Both insects bore into the leaf stalk just below the leaf blade. The leaf stalk shrivels, turns black, and the leaf blade falls off. The leaf drop may appear heavy but serious injury to a healthy tree is rare.

Aphids infest maples, usually Norway Maple, and may be numerous at times. High populations can cause leaf drop. Another sign of heavy aphid infestation is honey dew on lower leaves and objects beneath the tree. Aphids are controlled by spraying or they may be left alone. If not sprayed, predatory insects will usually bring the aphid population under control.

Scales are an occasional problem on maples. Perhaps the most common is cottony maple scale. The insect forms a cottony mass on the lower sides of branches. Scales are controlled with horticultural oil sprays. Scales may also be controlled with other well-timed sprays to kill the crawlers.

If borers become a problem it is an indication the tree is not growing well. Controlling borers involves keeping trees healthy. Chemical controls of existing infestations are more difficult. Proper control involves

identification of the borer infesting the tree then applying insecticides at the proper time.

Diseases

Verticillium wilt symptoms are wilting and death of branches. Infected sapwood will be stained a dark or olive green but staining can't always be found. If staining can not be found do not assume the problem is not verticillium wilt. Severely infected trees probably can't be saved. Lightly infected trees showing only a few wilted branches may be pulled through. Fertilize and prune lightly infected trees. This treatment will not cure the problem but may allow the tree to outgrow the infection. Girdling roots will cause symptoms which mimic verticillium wilt.

Tar spot and a variety of leaf spots cause some concern among homeowners but are rarely serious enough for control.