



Crataegus laevigata 'Crimson Cloud' 'Crimson Cloud' English Hawthorn¹

Edward F. Gilman and Dennis G. Watson²

INTRODUCTION

Crimson Cloud (also known as 'Superba') English Hawthorn grows rapidly in a pyramidal form to about 20 feet, then the crown expands to become oval or irregular (Fig. 1). The tree tolerates most soils, growing well in clay, but prefers heavy, dry loam. The main ornamental feature is white and red flowers borne in spring which together give the tree a deep pink color. Fruits are red and quite showy but do not cover the tree. Though quite ornamental, Hawthorns are susceptible to insect and disease problems. Branching habit is decidedly drooping and care should be given when locating this tree near pedestrian or vehicular traffic.

GENERAL INFORMATION

Scientific name: *Crataegus laevigata* 'Crimson Cloud'

Pronunciation: kruh-TEE-gus lee-vih-GAY-tuh

Common name(s): 'Crimson Cloud' English Hawthorn

Family: *Rosaceae*

USDA hardiness zones: 4B through 8 (Fig. 2)

Origin: not native to North America

Uses: Bonsai; espalier; wide tree lawns (>6 feet wide); medium-sized tree lawns (4-6 feet wide); recommended for buffer strips around parking lots or for median strip plantings in the highway; reclamation plant; screen; narrow tree lawns (3-4 feet wide); specimen; sidewalk cutout (tree pit); residential street tree; tree has been successfully grown in urban areas where air pollution, poor drainage, compacted soil, and/or drought are common




Figure 1. Middle-aged 'Crimson Cloud' English Hawthorn.

Availability: grown in small quantities by a small number of nurseries

DESCRIPTION

Height: 20 to 25 feet

Spread: 15 to 25 feet

Crown uniformity: irregular outline or silhouette

Crown shape: oval; pyramidal

Crown density: moderate

Growth rate: medium

Texture: fine

1. This document is adapted from Fact Sheet ST-211, a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Publication date: November 1993.
2. Edward F. Gilman, associate professor, Environmental Horticulture Department; Dennis G. Watson, associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.



Figure 2. Shaded area represents potential planting range.

Foliage

Leaf arrangement: alternate (Fig. 3)
Leaf type: simple
Leaf margin: lobed; serrate
Leaf shape: obovate; ovate
Leaf venation: pinnate
Leaf type and persistence: deciduous
Leaf blade length: less than 2 inches
Leaf color: green
Fall color: no fall color change
Fall characteristic: not showy

Flower

Flower color: lavender; red
Flower characteristics: showy; spring flowering

Fruit

Fruit shape: round
Fruit length: < .5 inch
Fruit covering: fleshy
Fruit color: red
Fruit characteristics: attracts birds; no significant litter problem; persistent on the tree; showy

Trunk and Branches

Trunk/bark/branches: droop as the tree grows, and will require pruning for vehicular or pedestrian clearance beneath the canopy; routinely grown with, or trainable to be grown with, multiple trunks; not particularly showy; tree wants to grow with several trunks but can be trained to grow with a single trunk; thorns are present on the trunk or branches
Pruning requirement: requires pruning to develop strong structure
Breakage: resistant
Current year twig color: brown
Current year twig thickness: medium

Culture

Light requirement: tree grows in full sun
Soil tolerances: clay; loam; sand; acidic; occasionally wet; alkaline; well-drained
Drought tolerance: high
Aerosol salt tolerance: low
Soil salt tolerance: poor




Figure 3. Foliage of 'Crimson Cloud' English Hawthorn.

Other

Roots: surface roots are usually not a problem

Winter interest: tree has winter interest due to unusual form, nice persistent fruits, showy winter trunk, or winter flowers

Outstanding tree: not particularly outstanding

Invasive potential: little, if any, potential at this time

Verticillium wilt susceptibility: not known to be susceptible

Pest resistance: long-term health usually not affected by pests

USE AND MANAGEMENT

The tree casts heavy shade, and turf can not be grown underneath, if lower branches are left on the trunk. Persistent lower branches add to the ornamental characteristic of the tree and it makes a nice specimen in a lawn for all seasons when left unpruned. When lower branches are removed in the nursery, this tough tree can be used as a street or parking lot tree where overhead space is limited by powerlines or other features. Be sure to order single-trunked trees clear of lower branches when planting along a street to provide for adequate clearance. Branches on the species bear sharp thorns which can inflict pain as they meet flesh, but this cultivar is almost thornless. Grows well in tree pits and other confined soil spaces. Plant in the open with good air circulation to help reduce disease, since leaf diseases can be common.

A number of other cultivars are offered in garden centers: 'Flore-Plena' - double, white flowers; 'Gireoudii' - foliage variegated, white flowers, large red fruit; 'Masekii' - pale rose, double flowers; 'Paulii' ('Paul's Scarlet') - deep pink, double flowers; 'Pendula' - weeping growth habit; 'Punica' - single, pink flowers; 'Rosea' - pink flowers; 'Stricta' - upright growth habit.

Pests

Aphids on small trees can be partially controlled with strong sprays of water from a garden hose, if the colony is in the lower branches. Sometimes the aphids themselves are not seen but the distorted growth, honeydew on the leaves, and sooty mold growing on the honeydew are obvious.

Borer attacks may be prevented if the trees are kept in good vigor with regular fertilization.

Leaf miners symptoms are brown blotches on the leaves.

Lace bugs can be a serious, though occasional, problem. The insect feeding on the undersides of the leaves causes chlorotic flecks on the upper leaf surfaces. The lower sides of the leaves are covered with small, brown, sticky flecks.

The pear slug skeletonizes Hawthorn leaves and these sawfly larvae have a slimy appearance. A few insects can be washed off with a garden hose.

Tent caterpillar nests can be pruned out while still small. Sprays of *Bacillus thuringiensis* may be used. Do not burn nests while the nests are in the tree. The injury from the fire may exceed that caused by the insects.

Scales may be controlled with horticultural oil sprays.

Spider mites are so small they can cause much foliage discoloration before being detected.

Diseases

Fire blight: This disease can be severe in some parts of the country. The first noticeable symptom of fire blight is the browning of branch tips. The tips appear to be burned or scorched and the dead, brown leaves droop but hang on the tree. Cankers form and the bacteria is washed farther down the branch by rain.

The bacteria, *Erwinia amylovora*, are spread from diseased to healthy twigs by rain, bees, and other mechanical means. There is no satisfactory chemical control. The disease is less of a problem if trees are not located near apple or pear orchards. Prune out blighted branch tips by cutting a foot or two beyond the diseased wood. Over-fertilizing with nitrogen fertilizer may increase tree susceptibility to fire blight.

Leaf blight attacks most Hawthorns but especially English Hawthorn. 'Crimson Cloud' is reported to be resistant. The symptoms are small reddish brown spots on the leaves which may run together. Infected leaves drop in August and severely infected trees may be completely bare.

Cedar Hawthorn rust causes orange or rust colored spots on the leaves leading to early defoliation. The fruits and twigs are also attacked. Juniper is an alternate host. Cedar-quince rust attacks fruits. Washington, Lavelle and Cockspur Hawthorn are resistant to rust diseases.

Scab causes leaf spotting and defoliation. The fruit have black raised spots on them.

Powdery mildew causes a white powdery growth on the leaves.