



Juniperus virginiana 'Pendula' 'Pendula' Eastern Redcedar¹

Edward F. Gilman and Dennis G. Watson²

INTRODUCTION

This cultivar of Redcedar is an evergreen growing 35 to 45 feet tall in a broad, oval form and spreading 15 to 25 feet when given a sunny location (Fig. 1). Branch tips droop toward the ground lending a weeping habit to the tree. This makes it a good candidate for a specimen to accent an area of the landscape. It develops a brownish tint in winter in the north and is sometimes used in windbreaks or screens. The fruit is a blue berry on female trees and is ornamental when produced in quantity. Birds devour the fruit and 'plant' it along farm fences and in old abandoned fields. Some botanists do not separate *Juniperus virginiana* from *Juniperus silicicola*.

GENERAL INFORMATION

Scientific name: *Juniperus virginiana* 'Pendula'

Pronunciation: joo-NIP-er-us ver-jin-ee-AY-nuh

Common name(s): 'Pendula' Eastern Redcedar

Family: Cupressaceae

USDA hardiness zones: 3 through 9 (Fig. 2)

Origin: native to North America

Uses: Bonsai; recommended for buffer strips around parking lots or for median strip plantings in the highway; reclamation plant; screen; specimen; tree has been successfully grown in urban areas where air pollution, poor drainage, compacted soil, and/or drought are common

Availability: somewhat available, may have to go out of the region to find the tree



Figure 1. Young 'Pendula' Eastern Redcedar.

DESCRIPTION

Height: 35 to 45 feet

Spread: 15 to 25 feet

Crown uniformity: symmetrical canopy with a regular (or smooth) outline, and individuals have more

1. This document is adapted from Fact Sheet ST-333, a series of the Environmental Horticulture Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Publication date: November 1993.
2. Edward F. Gilman, associate professor, Environmental Horticulture Department; Dennis G. Watson, associate professor, Agricultural Engineering Department, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.



Figure 2. Shaded area represents potential planting range.

or less identical crown forms

Crown shape: columnar; pyramidal; weeping

Crown density: moderate

Growth rate: fast

Texture: fine

Foliage

Leaf arrangement: opposite/subopposite; whorled

(Fig. 3)

Leaf type: simple

Leaf margin: entire; terminal spine

Leaf shape: awl-like; scale-like

Leaf venation: none, or difficult to see

Leaf type and persistence: evergreen

Leaf blade length: less than 2 inches

Leaf color: green

Fall color: no fall color change

Fall characteristic: not showy

Flower

Flower color: green; yellow

Flower characteristics: inconspicuous and not showy

Fruit

Fruit shape: round

Fruit length: < .5 inch

Fruit covering: fleshy

Fruit color: blue; purple

Fruit characteristics: attracts birds; no significant litter problem; persistent on the tree; showy

Trunk and Branches

Trunk/bark/branches: droop as the tree grows, and will require pruning for vehicular or pedestrian clearance beneath the canopy; showy trunk; should be grown with a single leader; no thorns

Pruning requirement: requires pruning to develop strong structure

Breakage: susceptible to breakage either at the crotch due to poor collar formation, or the wood itself is weak and tends to break

Current year twig color: brown; green

Current year twig thickness: thin

Wood specific gravity: 0.47




Figure 3. Foliage of 'Pendula' Eastern Redcedar.

Culture

Light requirement: tree grows in part shade/part sun;
tree grows in full sun

Soil tolerances: clay; loam; sand; acidic; alkaline;
well-drained

Drought tolerance: high

Aerosol salt tolerance: high

Other

Roots: surface roots are usually not a problem

Winter interest: no special winter interest

Outstanding tree: not particularly outstanding

Invasive potential: little, if any, potential at this time

Verticillium wilt susceptibility: not known to be
susceptible

Pest resistance: no pests are normally seen on the
tree

USE AND MANAGEMENT

The dense growth and attractive foliage make Eastern Redcedar a favorite for windbreaks, screens, and wildlife-cover for large-scale landscapes. Its high salt-tolerance makes it ideal for seaside locations. Redcedar can make a nice Christmas tree, and the fragrant wood is popular for repelling insects.

Planted in full sun or partial shade, Eastern Redcedar will easily grow on a variety of soils, including clay, but will not do well on soils kept continually moist. Growth will be poor in landscapes which are over-irrigated. Plants are difficult to transplant due to a coarse root system, except when quite small. Water until well-established and then forget about the tree. It performs admirably with no care, even on alkaline soil and along the coast. Usually insects and diseases are not a problem if grown in the full sun. There may be local restrictions on planting this tree near apple orchards because it is the alternate host for cedar-apple rust.

Some nurseries carry a cultivar or two of Redcedar.

Other cultivars include: 'Burkii' - pyramidal, blue foliage, 15 to 25 feet tall; 'Canaertii' - compact, pyramidal, good fruit production, fairly common in Texas; 'Hillspire' - (cupressifolia) - good green color; 'Elegantissima' - Goldtip Redcedar - branchlets with yellow tips, less than 20 feet tall; 'Filifera' - pyramidal, branchlets divided, foliage gray green; 'Glauca' - Silver Redcedar - narrow, columnar, 15 to 20 feet tall, silvery blue foliage especially in spring; 'Ketlerii' is commonly available in the mid-west, is more open with spaces between branches at the top of the tree, pyramidal; 'Manhattan Blue' - compact, 20 feet tall, pyramidal, foliage bluish green; 'Pyramidalis Dundee' - pyramidal, purplish green in winter; 'Skyrocket' - silver-blue foliage, narrow columnar form.

Pests

Usually none are serious.

Bagworm caterpillars occasionally web foliage and debris together to make bags up to two inches long. The insects live in the bags and emerge to feed on the foliage. Use sprays of *Bacillus thuringiensis*. The insects can also be picked off the plants by hand.

Juniper scale causes yellowed needles, and infected branches fail to produce new growth. The scale is round and at first white, later turning gray or black.

The Juniper webworm webs twigs and needles together, causing them to brown and die. The larva is 1/2-inch-long and is brown with darker stripes. The larvae are often in the densest part of the plant and can go unnoticed.

Mites cause stippled and bronzed foliage.

Diseases

Twig blights cause death and browning of twig tips. The diseases may progress down the stem killing the whole branch. Small lesions may be seen at the base of dead tissue. Prune out dead branch tips. Dieback from *Kabatina* blight appears in early spring, from *Phomopsis* in summer.

Three rust diseases seen most often are cedar-apple rust, hawthorn rust, and quince rust. The most common is cedar-apple rust. On Juniper the first two diseases form galls and orange jelly-like horns in spring. The horns are most likely to form following periods of rainy, warm weather. Spores formed in the horns infect the alternate host. The diseases are more serious on the alternate host than Juniper. There may be local restrictions on planting this tree near apple orchards because it is the alternate host for cedar-apple rust. A separation of a few hundred yards may help avoid the disease. Prune out the spore horns when seen in the spring. Do not plant near hawthorns, apples, or crabapples.

Junipers are not tolerant of ice coatings. Expect dieback when Junipers are covered with ice for several days. Removing the ice is impractical.